

Asymptotic typicality degrees of properties over finite structures

Athanassios Tzouvaras

Department of Mathematics, Aristotle University of Thessaloniki, 541 24 Thessaloniki, Greece

Abstract. In previous work we defined and studied a notion of typicality, originated with B. Russell, for properties and objects in the context of general infinite first-order structures. In this paper we consider this notion in the context of finite structures. In particular we define the typicality degree of a property $\varphi(x)$ over finite L -structures, for a language L , as the limit of the probability of $\varphi(x)$ to be typical in an arbitrary L -structure \mathcal{M} of cardinality n , when n goes to infinity. This poses the question whether the 0-1 law holds for typicality degrees for certain kinds of languages. One of the results of the paper is that, in contrast to the classical well-known fact that the 0-1 law holds for the sentences of every relational language, the 0-1 law fails for degrees of properties of relational languages containing unary predicates. On the other hand it is shown that the 0-1 law holds for degrees of some basic properties of graphs, and this gives rise to the conjecture that the 0-1 law holds for relational languages without unary predicates. Another theme is the *neutrality* degree of a property $\varphi(x)$ (i.e., the fraction of L -structures in which neither φ nor $\neg\varphi$ is typical), and in particular the *regular* properties (i.e., those with limit neutrality degree 0). All properties we dealt with, either of a relational or a functional language, are shown to be regular, but the question whether *every* such property is regular is open.

1 Typicality à la Russell

In [9], we set out an investigation of a notion of typicality which is originated with Bertrand Russell. Specifically in [7, p. 89], Russell defines a typical Englishman to be one “who possesses all the properties possessed by a majority of Englishmen.” The notion seems captivating in its simplicity and naturalness, but in order to be formally defined one has to distinguish between properties of an object language and properties of the metalanguage, else typicality itself would be one of the properties we have to check about an Englishman, and thus circularity arises. Once we make the aforementioned distinction using a first-order language L , given any L -structure $\mathcal{M} = \langle M, \dots \rangle$ we can define typical elements of M in the spirit of Russell, provided we first define what a typical property is. Given a formula $\varphi(x)$ of L without parameters, let $\varphi(\mathcal{M})$ denote the extension of $\varphi(x)$ in \mathcal{M} , i.e., $\varphi(\mathcal{M}) = \{a \in M : \mathcal{M} \models \varphi(a)\}$.

Definition 1. Let $\mathcal{M} = \langle M, \dots \rangle$ be an L -structure. A property $\varphi(x)$ of L is said to be *typical over* \mathcal{M} , if $|\varphi(\mathcal{M})| > |\neg\varphi(\mathcal{M})| = |M \setminus \varphi(\mathcal{M})|$. Then an element $a \in M$ is said to be *typical* if it satisfies every typical property over \mathcal{M} .

In [9], among other things, we established the existence of typical elements in many *infinite* structures. For example the standard structure of reals (or second-order arithmetic) contains $|\mathbb{R}|$ -many typical reals, while

only $<|\mathbb{R}|$ -many nontypical ones. (A variant of the same notion of typicality, adjusted to fit to the context of set theory and generating a new inner model of ZF, has appeared in [10].)

Instead, in the present paper we are interested only in typical *properties* (not in typical objects), and only over the *finite* structures of a (finite) language L . Specifically, we set out to study the *probabilities* for L -properties $\varphi(x)$, in one free variable, to be *typical* over arbitrary L -structures \mathcal{M} of cardinality n , and further to compute the limits of these probabilities, as n tends to infinity. This study parallels the classical results of finite model theory about the probabilities of *sentences* of L to be *true* over finite structures and the fundamental 0-1 law about these truth probabilities.

2 Typicality degrees of first-order properties over finite structures

2.1 Asymptotic truth probabilities and 0-1 laws

Typicality degrees of properties over finite structures and their asymptotic behavior are, in some sense, generalizations of the truth degrees (or truth probabilities) of *sentences*. So we need to recall first some definitions and notation about the latter, see for example [2, § 3], or the original paper [3]. The terminology and notation here is mostly that of [3].

Let L be a first-order language consisting of a finite set of (non-logical) relational symbols. For every n , let $\mathbf{S}_n(L)$ be the set of L -structures $\mathcal{M} = \langle M, \dots \rangle$ with $|M| = n$, or simply $M = \{1, 2, \dots, n\}$. (In doing so, of course, we tacitly use the fact that if any two models, finite or infinite, $\mathcal{M} = \langle M, \dots \rangle$, $\mathcal{N} = \langle N, \dots \rangle$, differ *only in the nature* of elements of M, N , while $|M| = |N|$, then they are isomorphic, so mathematically identical). For every L -sentence φ , let $\text{Mod}_n(\varphi)$ be the subset of structures of $\mathbf{S}_n(L)$ which satisfy φ . Let also

$$\mu_n(\varphi) = \frac{|\text{Mod}_n(\varphi)|}{|\mathbf{S}_n(L)|}, \quad \text{and} \quad \mu(\varphi) = \lim_{n \rightarrow \infty} \mu_n(\varphi),$$

if this limit exists. Given a class Φ of L -sentences we say that Φ *satisfies the 0-1 law* if for every $\varphi \in \Phi$, $\mu(\varphi) = 0$ or $\mu(\varphi) = 1$. The following is a fundamental result of finite model theory. The following theorem is independently due to Fagin and Glebskii et al. [3; 4].

Theorem 2. (0-1 law for first-order logic) If L is a first-order language with no function or constant symbols, then the set of sentences of L satisfies the 0-1 law.

Nevertheless, Theorem 2 fails when L contains function symbols. The following is the standard example used to prove this failure (see [3, § 4] and [2, Example 3.1.1]).

Example 3. Let $L = \{F\}$, where F is a unary function symbol. If φ is the L -sentence $\varphi := \forall x(F(x) \neq x)$, then $\mu(\varphi) = e^{-1}$, thus the 0-1 law fails in general for the sentences of L .

Proof. Observe that for $n \geq 1$, the number of structures $\mathcal{M} = \langle M, f \rangle \in \mathbf{S}_n(L)$ that satisfy $\varphi := (\forall x)(F(x) \neq x)$ is just the number of functions $f : M \rightarrow M$, $|M| = n$, such that $f(x) \neq x$ for every $x \in M$. This number is $(n-1)^n$ (since $f(x)$ may take independently for each x , $n-1$ possible values). On the other hand, $|\mathbf{S}_n(L)| = n^n$. Therefore, $\mu_n(\varphi) = (1 - 1/n)^n$, and hence $\lim_{n \rightarrow \infty} \mu_n(\varphi) = e^{-1}$. \square

2.2 Typicality degrees of properties

Let us first elaborate a bit on the general Definition 1. Recall that we denote by $\varphi(\mathcal{M})$ the extension of φ is \mathcal{M} , i.e., $\varphi(\mathcal{M}) = \{a \in M : \mathcal{M} \models \varphi(a)\}$. When we deal with typicality of *elements* of a structure, we naturally distinguish them into just typical and non-typical, but when we deal with typicality of *properties*, especially over finite structures, we should distinguish them into three kinds, according to the size of their extension.

Definition 4. Let \mathcal{M} be an L -structure and let $\varphi(x)$ be a property of L (without parameters). We say that:

- (1) $\varphi(x)$ is *typical* for \mathcal{M} , if $|\varphi(\mathcal{M})| > |\neg\varphi(\mathcal{M})|$.
- (2) $\varphi(x)$ is *atypical* for \mathcal{M} , if $|\varphi(\mathcal{M})| < |\neg\varphi(\mathcal{M})|$.
- (3) $\varphi(x)$ is *neutral* for \mathcal{M} , if $|\varphi(\mathcal{M})| = |\neg\varphi(\mathcal{M})|$ (i.e., if neither $\varphi(x)$ nor $\neg\varphi(x)$ is typical).

The above distinction of properties is valid for all structures, infinite and finite, but is particularly useful when dealing with finite structures. If we apply the preceding definition to a structure \mathcal{M} of $\mathbf{S}_n(L)$, then $\varphi(x)$ is typical, atypical and neutral for \mathcal{M} , if and only if $|\varphi(\mathcal{M})| > n/2$, $|\varphi(\mathcal{M})| < n/2$ and $|\varphi(\mathcal{M})| = n/2$, respectively, the latter case of course being possible only for even n . To simplify terminology, henceforth we shall not refer to *atypical* $\varphi(x)$ but to *typical* $\neg\varphi(x)$ instead, since for every φ and $\mathcal{M} \in \mathbf{S}_n(L)$, $|\varphi(\mathcal{M})| < n/2$ iff $|\neg\varphi(\mathcal{M})| > n/2$, that is, $\varphi(x)$ is atypical for \mathcal{M} iff $\neg\varphi(x)$ is typical for \mathcal{M} . Let us also fix for every L and n the following subclasses of $\mathbf{S}_n(L)$.

$$\mathbf{S}_n(\varphi : \text{typ}) = \{\mathcal{M} \in \mathbf{S}_n(L) : \varphi(x) \text{ is typical for } \mathcal{M}\} = \{\mathcal{M} : |\varphi(\mathcal{M})| > n/2\},$$

$$\mathbf{S}_n(\varphi : \text{ntr}) = \{\mathcal{M} \in \mathbf{S}_n(L) : \varphi(x) \text{ is neutral for } \mathcal{M}\} = \{\mathcal{M} : |\varphi(\mathcal{M})| = n/2\}.$$

The second of the above classes is nonempty only for even n , so for each property $\varphi(x)$, $\mathbf{S}_n(L)$ splits as follows for odd and even n :

$$\mathbf{S}_{2n+1}(L) = \mathbf{S}_{2n+1}(\varphi : \text{typ}) \cup \mathbf{S}_{2n+1}(\neg\varphi : \text{typ}), \quad (1)$$

$$\mathbf{S}_{2n}(L) = \mathbf{S}_{2n}(\varphi : \text{typ}) \cup \mathbf{S}_{2n}(\neg\varphi : \text{typ}) \cup \mathbf{S}_{2n}(\varphi : \text{ntr}). \quad (2)$$

Then, by analogy with the probabilities $\mu_n(\varphi)$, and asymptotic probability $\mu(\varphi) = \lim_{n \rightarrow \infty} \mu_n(\varphi)$ for the truth of L -sentences referred to in Section 2.1 above, we can naturally define the corresponding probabilities for a property $\varphi(x)$ to be typical or neutral over an arbitrary structure $\mathcal{M} \in \mathbf{S}_n(L)$. Specifically, for each n , we set

$$d_n(\varphi : \text{typ}) = \frac{|\mathbf{S}_n(\varphi : \text{typ})|}{|\mathbf{S}_n(L)|}, \quad d_{2n}(\varphi : \text{ntr}) = \frac{|\mathbf{S}_{2n}(\varphi : \text{ntr})|}{|\mathbf{S}_{2n}(L)|},$$

$$d(\varphi : \text{typ}) = \lim_{n \rightarrow \infty} d_n(\varphi : \text{typ}), \quad d(\varphi : \text{ntr}) = \lim_{n \rightarrow \infty} d_{2n}(\varphi : \text{ntr}),$$

whenever these limits exist. Here, $d_n(\varphi : \text{typ})$ and $d_{2n}(\varphi : \text{ntr})$ are the n -*typicality degree* and n -*neutrality degree* of $\varphi(x)$, respectively, while $d(\varphi : \text{typ})$ and $d(\varphi : \text{ntr})$ are the corresponding *asymptotic degrees*. Here are some first consequences of the definitions.

Fact 5.

- (i) If $\vdash \varphi(x) \rightarrow \psi(x)$, then $d_n(\varphi : \text{typ}) \leq d_n(\psi : \text{typ})$, for all $n \geq 1$. Therefore, if $d(\psi : \text{typ}) = 0$, then $d(\varphi : \text{typ}) = 0$ too.
- (ii) If $\vdash \varphi(x) \leftrightarrow \psi(x)$, then $d_n(\varphi : \text{typ}) = d_n(\psi : \text{typ})$, and also $d_{2n}(\varphi : \text{ntr}) = d_{2n}(\psi : \text{ntr})$ for all $n \geq 1$.
- (iii) For all $\varphi(x)$ and n , $d_{2n}(\varphi : \text{ntr}) = d_{2n}(\neg\varphi : \text{ntr})$.
- (iv) If $d(\varphi : \text{typ}) = a > 0$ (resp. $d(\varphi : \text{ntr}) = a > 0$), then the set $\{n : \mathbf{S}_n(\varphi : \text{typ}) \neq \emptyset\}$ (resp. $\{n : \mathbf{S}_{2n}(\varphi : \text{ntr}) \neq \emptyset\}$) is cofinite.

Proof. For (i) and (ii) just note that if $\vdash \varphi(x) \rightarrow \psi(x)$, then for every structure \mathcal{M} , $\varphi(\mathcal{M}) \subseteq \psi(\mathcal{M})$, hence $|\varphi(\mathcal{M})| \leq |\psi(\mathcal{M})|$, so

$$\mathcal{M} \in \mathbf{S}_n(\varphi : \text{typ}) \Leftrightarrow |\varphi(\mathcal{M})| > n/2 \Rightarrow |\psi(\mathcal{M})| > n/2 \Leftrightarrow \mathcal{M} \in \mathbf{S}_n(\psi : \text{typ}),$$

therefore, $\mathbf{S}_n(\varphi : \text{typ}) \subseteq \mathbf{S}_n(\psi : \text{typ})$. Moreover $\vdash \varphi(x) \leftrightarrow \psi(x)$ implies $\mathbf{S}_n(\varphi : \text{typ}) = \mathbf{S}_n(\psi : \text{typ})$ and also $\mathbf{S}_{2n}(\varphi : \text{ntr}) = \mathbf{S}_{2n}(\psi : \text{ntr})$, for every n .

- (iii) If $|M| = 2n$, then for every $\varphi(x)$, obviously $|\varphi(\mathcal{M})| = n$ iff $|\neg\varphi(\mathcal{M})| = n$, so $\mathbf{S}_{2n}(\varphi : \text{ntr}) = \mathbf{S}_{2n}(\neg\varphi : \text{ntr})$.

(iv) Let $d(\varphi : \text{typ}) = a > 0$. If we pick some $0 < \varepsilon < a$, then clearly there is n_0 such that for all $n \geq n_0$, $d_n(\varphi : \text{typ}) > a - \varepsilon$. Since

$$d_n(\varphi : \text{typ}) = \frac{|\mathbf{S}_n(\varphi : \text{typ})|}{|\mathbf{S}_n(L)|},$$

it follows that for all $n \geq n_0$ $|\mathbf{S}_n(\varphi : \text{typ})| \neq 0$, and hence $\mathbf{S}_n(\varphi : \text{typ}) \neq \emptyset$. The claim for $\{n : \mathbf{S}_{2n}(\varphi : \text{ntr}) \neq \emptyset\}$ is shown similarly. \square

By the definition of $\mu(\varphi)$ in Section 2.1, it follows immediately that for any language L and any L -sentence φ , if $\mu(\varphi)$ exists, then so does $\mu(\neg\varphi)$ and $\mu(\neg\varphi) = 1 - \mu(\varphi)$. What about typicality degrees? Is it true that $d(\neg\varphi : \text{typ}) = 1 - d(\varphi : \text{typ})$ whenever $d(\varphi : \text{typ})$ exists, for any property $\varphi(x)$? The question is eventually open and the reason is the limit $\lim_{n \rightarrow \infty} d_{2n}(\varphi : \text{ntr})$. Namely, while by (1) $d_{2n+1}(\varphi : \text{typ}) + d_{2n+1}(\neg\varphi : \text{typ}) = 1$, (2) implies that $d_{2n}(\varphi : \text{typ}) + d_{2n}(\neg\varphi : \text{typ}) + d_{2n}(\varphi : \text{ntr}) = 1$. Thus, in the second case we have

$$\lim_{n \rightarrow \infty} d_{2n}(\neg\varphi : \text{typ}) = 1 - \lim_{n \rightarrow \infty} d_{2n}(\varphi : \text{typ}) - \lim_{n \rightarrow \infty} d_{2n}(\varphi : \text{ntr}),$$

and in order to infer that

$$\lim_{n \rightarrow \infty} d_{2n}(\neg\varphi : \text{typ}) = 1 - \lim_{n \rightarrow \infty} d_{2n}(\varphi : \text{typ}),$$

we must establish that $\lim_{n \rightarrow \infty} d_{2n}(\varphi : \text{ntr}) = 0$. We don't know if this is the case for every property $\varphi(x)$ of every language. So we shall give a name to properties satisfying this interesting and convenient condition.

3 Regularity of properties

Definition 6. A property $\varphi(x)$ of L is said to be *regular* if $d(\varphi : \text{ntr}) = 0$.

Fact 7.

- (i) $\varphi(x)$ is regular if and only if $\neg\varphi(x)$ is regular.
- (ii) If $d(\varphi : \text{typ})$ exists, then so does $\lim_{n \rightarrow \infty} d_{2n+1}(\neg\varphi : \text{typ})$ and $\lim_{n \rightarrow \infty} d_{2n+1}(\neg\varphi : \text{typ}) = 1 - d(\varphi : \text{typ})$.
- (iii) If $\varphi(x)$ is regular, then also $\lim_{n \rightarrow \infty} d_{2n}(\neg\varphi : \text{typ}) = 1 - d(\varphi : \text{typ})$, and thus, $d(\neg\varphi : \text{typ}) = 1 - d(\varphi : \text{typ})$.

Proof. (i) By Fact 5 (iii), for every n , $d_{2n}(\varphi : \text{ntr}) = d_{2n}(\neg\varphi : \text{ntr})$, therefore $d(\varphi : \text{ntr}) = 0$ if and only if $d(\neg\varphi : \text{ntr}) = 0$.

- (ii) If $d(\varphi : \text{typ}) = a$, then also $\lim_{n \rightarrow \infty} d_{2n+1}(\varphi : \text{typ}) = a$, thus by (1), $\lim_{n \rightarrow \infty} d_{2n+1}(\neg\varphi : \text{typ}) = 1 - a$.
- (iii) If $d(\varphi : \text{typ}) = a$ and $\varphi(x)$ is regular, then $\lim_{n \rightarrow \infty} d_{2n}(\varphi : \text{ntr}) = 0$, so by (2)

$$\lim_{n \rightarrow \infty} d_{2n}(\neg\varphi : \text{typ}) = 1 - a - \lim_{n \rightarrow \infty} d_{2n}(\varphi : \text{ntr}) = 1 - a. \quad \square$$

All specific properties we treat below are regular. So it is natural to ask whether every property is regular. The question is open for general languages. In the next subsection we show that it is true for a large class of properties of the language $L = \{U_1, \dots, U_k\}$ which consists of an arbitrary number of unary predicates.

3.1 Languages with finitely many unary predicates

Let $L = \{U_1, \dots, U_k\}$ be a language with k unary predicates. For each $i \in \{1, \dots, k\}$, let $U_i^1(x) = U_i(x)$ and $U_i^0(x) = \neg U_i(x)$. Then given a function $e \in \{0, 1\}^k$, we set $\varphi_e(x) = U_1^{e(1)}(x) \wedge U_2^{e(2)}(x) \wedge \dots \wedge U_k^{e(k)}(x)$. The properties $\varphi_e(x)$, for $e \in \{0, 1\}^k$, form the 2^k atoms of the (syntactic) Boolean algebra \mathcal{B}_{prop} generated by the properties $U_i(x)$, $i \in \{1, \dots, k\}$, and any two distinct atoms $\varphi_{e_1}(x)$, $\varphi_{e_2}(x)$ are mutually inconsistent, i.e., $\varphi_{e_1}(x) \wedge \varphi_{e_2}(x) \vdash \perp$. Besides each of the 2^{2^k} elements of \mathcal{B}_{prop} has the form $\varphi_E(x) = \bigvee_{e \in E} \varphi_e(x)$, for some $E \subseteq \{0, 1\}^k$.

We shall generalize the class of formulas $\varphi_e(x)$ defined above, by relaxing the condition that for every $i \leq k$, either $U_i(x)$ or $\neg U_i(x)$ must be a conjunct of φ_e . Namely for any p such that $1 \leq p \leq k$, a p -subsequence of $\langle 1, \dots, k \rangle$ is a p -tuple $\langle i_1, \dots, i_p \rangle$ such that $1 \leq i_1 < i_2 < \dots < i_p \leq k$. Given a p -subsequence $\bar{s} = \langle i_1, \dots, i_p \rangle$ and an $e \in \{0, 1\}^k$, let $\varphi_{\bar{s},e}(x) = U_{i_1}^{e(i_1)}(x) \wedge \dots \wedge U_{i_p}^{e(i_p)}(x)$. We refer to formulas $\varphi_{\bar{s},e}(x)$ as *basic formulas* of L . The main result of this subsection is that every basic formula of L is regular.

Given an L -structure $\mathcal{M} = \langle M, W_1, \dots, W_k \rangle$, let $W_i^1 = W_i$ and $W_i^0 = M \setminus W_i$. For every $e \in \{0, 1\}^k$, let $X_e = W_1^{e(1)} \cap W_2^{e(2)} \cap \dots \cap W_k^{e(k)}$. Clearly the sets X_e , for $e \in \{0, 1\}^k$, are pairwise disjoint, but their difference from $\varphi_e(x)$ is that not all of them need to be nonempty. Let $\mathcal{B}_{set} \subseteq \mathcal{P}(M)$ be the Boolean algebra generated by the sets W_i , $i \in \{1, \dots, k\}$ with at most 2^k atoms. As before for each $E \subseteq \{0, 1\}^k$, we set $X_E = \bigcup_{e \in E} X_e$. Further for any p -subsequence $\bar{s} = \langle i_1, \dots, i_p \rangle$ of $\langle 1, \dots, k \rangle$, with $1 \leq p \leq k$, and any $e \in \{0, 1\}^k$, we write $X_{\bar{s},e} = W_{i_1}^{e(i_1)} \cap \dots \cap W_{i_p}^{e(i_p)}$. Finally each set X_E is defined in \mathcal{M} by the property $\varphi_E(x)$, i.e., $X_E = \varphi_E(\mathcal{M})$, and each $X_{\bar{s},e}$ is defined by $\varphi_{\bar{s},e}(x)$, i.e., $\varphi_{\bar{s},e}(\mathcal{M}) = X_{\bar{s},e}$.

Lemma 8. Given an L -structure $\mathcal{M} = \langle M, W_1, \dots, W_k \rangle$ as above, the definable (without parameters) subsets of M are exactly the sets X_E . Therefore, every L -property $\varphi(x)$ is equivalent over \mathcal{M} to a $\varphi_E(x)$, for some $E \subseteq \{0, 1\}^k$.

Proof. The proof is easy and based on the fact that for any two elements $a \neq b \in M$, there is an automorphism $f : M \rightarrow M$ which interchanges a and b . \square

Let us notice here a result which is involved in all proofs of regularity of properties. Whenever we try to check the regularity of a property over a structure M with $2n$ elements, we shall necessarily deal with the number $\binom{2n}{n}$ which counts the subsets M having half of its elements. The numbers $\binom{2n}{n}$ are known as *central binomial coefficients* and several useful combinatorial facts are known about them (see, e.g., [8]). In particular the following upper bound is particularly helpful and will be used below.

Fact 9. For every $n \geq 1$, $\binom{2n}{n} \leq \frac{4^n}{\sqrt{\pi n}}$.

For $k \leq n$, we shall denote by $(n)_k$ the number of k -tuples of distinct elements chosen from a set of n elements. It is well known that

$$(n)_k = \frac{n!}{(n-k)!} = (n-k+1)(n-k+2) \cdots n.$$

Then $(n)_1 = n$, $(n)_n = n!$, $(n)_k = 0$ for $k > n$. In particular we set $(n)_0 = 1$. The numbers $(n)_m$ are called *falling factorials*. Notations n^m and $P(n, m)$ are often used in the bibliography instead of $(n)_m$.

Below we shall extensively employ the relation of *asymptotic equality* between functions $f, g : \mathbb{N} \rightarrow \mathbb{R}$, denoted $f(n) \sim g(n)$, as well as that of *asymptotic inequality* $f(n) \lesssim g(n)$. Both these relations are based on the well-known relation of *little-o notation* between functions f, g , denoted $f(n) = o(g(n))$ (see, e.g., [1]). For reader's convenience below we recall the basic definitions and properties of the aforementioned relations.

Firstly, for a given function $g(n)$, $og(n)$ denotes any function such that $\lim_{n \rightarrow \infty} \frac{og(n)}{g(n)} = 0$, i.e., $f(n) = og(n)$ iff $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$. Next, $f(n) \sim g(n)$ means $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 1$, or equivalently, $f(n) = g(n) + og(n)$. Similarly, $f(n) \lesssim g(n)$ means $f(n) \leq g(n) + o(g(n))$. The properties of \sim and \lesssim that we shall need below are the following, and they are either well-known or easily verified.

Fact 10.

- (i) The relation \sim is preserved by the usual operations, i.e., if $f_1 \sim g_1$ and $f_2 \sim g_2$, then $f_1 + f_2 \sim g_1 + g_2$, $f_1 \cdot f_2 \sim g_1 \cdot g_2$, and $\frac{f_1}{f_2} \sim \frac{g_1}{g_2}$.
- (ii) If $f(n) \sim g(n)$ and $\lim_{n \rightarrow \infty} f(n) = a \in \mathbb{R}$, then $\lim_{n \rightarrow \infty} g(n) = a$.
- (iii) If $f(n) \sim g(n)$ and $g(n) \leq h(n)$, then $f(n) \lesssim h(n)$.
- (iv) If $0 \leq f(n) \lesssim g(n)$ and $\lim_{n \rightarrow \infty} g(n) = 0$, then $\lim_{n \rightarrow \infty} f(n) = 0$.

Proof. For illustration we shall just sketch the proof of (ii). Let $f(n) \sim g(n)$ and $\lim_{n \rightarrow \infty} f(n) = a \in \mathbb{R}$. Then clearly both f, g are bounded, and let b be a bound for g , i.e., $\forall n |g(n)| \leq b$. We have $f(n) = g(n) + o(g(n))$, where $\lim_{n \rightarrow \infty} \frac{o(g(n))}{g(n)} = 0$. So for all n

$$|g(n) - a| \leq |g(n) - f(n)| + |f(n) - a| = |o(g(n))| + |f(n) - a|. \quad (3)$$

Fix some $\varepsilon > 0$. There is n_1 such that $\forall n \geq n_1 \left| \frac{o(g(n))}{g(n)} \right| \leq \frac{\varepsilon}{2b}$, hence

$$\forall n \geq n_1 |o(g(n))| \leq \frac{\varepsilon}{2b} |g(n)| \leq \frac{\varepsilon}{2}. \quad (4)$$

Also there is n_2 such that

$$\forall n \geq n_2 |f(n) - a| < \frac{\varepsilon}{2}. \quad (5)$$

If $n_0 = \max(n_1, n_2)$, (3), (4) and (5) yield $\forall n \geq n_0 |g(n) - a| < \varepsilon$. \square

For example, for any fixed k such that $1 \leq k < n$, $(n)_k$ is a polynomial in n of degree k with leading coefficient 1, so

$$(n)_k \sim n^k. \quad (6)$$

We shall apply relation (6) several times without explicit reference to that. We shall also make use of the following result.

Fact 11. $\sum_{k < n/2} \binom{n}{k} = \sum_{k > n/2} \binom{n}{k} \sim 2^{n-1}$.

Proof. The first equality is well-known. Moreover for every n , $\sum_{k=0}^n \binom{n}{k} = 2^n$. If n is odd,

$$\sum_{k=0}^n \binom{n}{k} = \sum_{k < n/2} \binom{n}{k} + \sum_{k > n/2} \binom{n}{k},$$

so

$$\sum_{k < n/2} \binom{n}{k} = \frac{2^n}{2} = 2^{n-1}.$$

If n is even and $n = 2m$, then

$$2 \left(\sum_{k < n/2} \binom{n}{k} \right) + \binom{2m}{m} = 2^n,$$

that is,

$$\sum_{k < n/2} \binom{n}{k} = \frac{1}{2} \left(2^n - \binom{2m}{m} \right) = 2^{n-1} - \frac{1}{2} \binom{2m}{m}.$$

Therefore,

$$\frac{\sum_{k < n/2} \binom{n}{k}}{2^{n-1}} = 1 - \frac{1}{2^{2m}} \binom{2m}{m}.$$

So it suffices to see that $\lim_{m \rightarrow \infty} \frac{1}{2^{2m}} \binom{2m}{m} = 0$. But this follows from Fact 9, since

$$\frac{1}{2^{2m}} \binom{2m}{m} \leq \frac{1}{2^{2m}} \frac{4^m}{\sqrt{\pi m}} = \frac{1}{\sqrt{\pi m}} \rightarrow_m 0. \quad \square$$

Before going on we shall make the following important clarification/simplification. According to the standard basic definition of first-order logic, an interpretation of a language $L = \{U_1, \dots, U_k\}$ with unary predicates in a set M , is any mapping $I : \{U_1, \dots, U_k\} \rightarrow \mathcal{P}(M)$, where the set $I(U_i) = W_i \subseteq M$ interprets U_i , i.e., is the extension of U_i , and is written also $W_i = U_i^{\mathcal{M}}$, where \mathcal{M} is the model on M created by this interpretation. This means that the class of models of L of cardinality n , $\mathbf{S}_n(L)$, has cardinality equal to the number of mappings $I : \{U_1, \dots, U_k\} \rightarrow \mathcal{P}(M)$, for $|M| = n$, i.e., $|\mathbf{S}_n(L)| = |\mathcal{P}(M)|^k = 2^{kn}$. Notice that such arbitrary interpretations $I : \{U_1, \dots, U_k\} \rightarrow \mathcal{P}(M)$ may assign the same extension to different predicates U_i of L , i.e., we may have $U_i^{\mathcal{M}} = U_j^{\mathcal{M}}$ for $i \neq j$.

This fact, however, increases significantly the combinatorial complexity that one meets when trying to compute properties like typicality or regularity of properties of the language L , which is the object of the present paper.

The complexity can be significantly reduced and the computations be much more accessible, if instead of arbitrary interpretations $I : \{U_1, \dots, U_k\} \rightarrow \mathcal{P}(M)$ of L , we had to deal only with *1-1 interpretations*, i.e., with models $\mathcal{M} = \langle W_1, \dots, W_k \rangle$ such that $W_i = U_i^{\mathcal{M}} \neq W_j = U_j^{\mathcal{M}}$, for all $i \neq j$. After all it is intuitively much more natural to assign different extensions $U_i^{\mathcal{M}} \neq U_j^{\mathcal{M}}$ to different predicates U_i, U_j . We refer to such models as *1-1 models*.

The class of 1-1 models, let us denote it $\mathbf{S}_n^{1-1}(L)$, is a subclass of $\mathbf{S}_n(L)$, however it is easy to see that $|\mathbf{S}_n^{1-1}(L)|$ is *asymptotically equal* to $|\mathbf{S}_n(L)|$, i.e., $|\mathbf{S}_n^{1-1}(L)| \sim |\mathbf{S}_n(L)|$. Indeed, $\mathbf{S}_n^{1-1}(L)$ is essentially the set of 1-1 mappings $I : \{U_1, \dots, U_k\} \rightarrow \mathcal{P}(M)$, and each such mapping can be identified with a k -tuple of distinct elements of $\mathcal{P}(M)$. Therefore, $|\mathbf{S}_n^{1-1}(L)|$ is equal to the number of all such k -tuples of $\mathcal{P}(M)$, i.e., $(2^n)_k$. But $(2^n)_k \sim 2^{kn}$ according to (6). So indeed $|\mathbf{S}_n^{1-1}(L)| \sim |\mathbf{S}_n(L)|$.

I am sure that the other classes of models which are involved in the computation of asymptotic typicality and regularity of properties of the language L with unary predicates, namely $\mathbf{S}_n(\varphi_{\bar{s},e} : \text{typ})$ and $\mathbf{S}_{2n}(\varphi_{\bar{s},e} : \text{ntr})$, remain asymptotically equal to $\mathbf{S}_n(\varphi_{\bar{s},e} : \text{typ})$ and $\mathbf{S}_{2n}(\varphi_{\bar{s},e} : \text{ntr})$, respectively, when relativized to the class $\mathbf{S}_n^{1-1}(L)$, i.e., $\mathbf{S}_n^{1-1}(\varphi_{\bar{s},e} : \text{typ}) \sim \mathbf{S}_n(\varphi_{\bar{s},e} : \text{typ})$, and $\mathbf{S}_{2n}^{1-1}(\varphi_{\bar{s},e} : \text{ntr}) \sim \mathbf{S}_{2n}(\varphi_{\bar{s},e} : \text{ntr})$. However the proof of this does not seem to be an easy task.

Nevertheless, below I shall choose to work with models of $\mathbf{S}_n^{1-1}(L)$ rather than $\mathbf{S}_n(L)$, because the computations with them are tractable and accessible to the reader. In the worst case that $\mathbf{S}_n^{1-1}(\varphi_{\bar{s},e} : \text{typ}) \not\sim \mathbf{S}_n(\varphi_{\bar{s},e} : \text{typ})$, and/or $\mathbf{S}_{2n}^{1-1}(\varphi_{\bar{s},e} : \text{ntr}) \not\sim \mathbf{S}_{2n}(\varphi_{\bar{s},e} : \text{ntr})$, the results can be simply considered as *partial results* referring only to a subclass of the class of all models.

After the above explanations we proceed to compute typicality and regularity, with the proviso that we deal with 1-1 models only. Since below there is no need to keep the notation $\mathbf{S}_n^{1-1}(L)$, we return to the notation $\mathbf{S}_n(L)$.

Theorem 12. Every basic property $\varphi_{\bar{s},e}(x)$ of $L = \{U_1, \dots, U_k\}$ is regular. In particular every property $\varphi_e(x)$, as well as every $U_i(x)$ and $\neg U_i(x)$, for $i = 1, \dots, k$, is regular.

Proof. Let us fix the ground set M of all $\mathcal{M} = \langle M, W_1, \dots, W_k \rangle \in \mathbf{S}_{2n}(L)$, i.e., $|M| = 2n$, and fix also a basic formula $\varphi_{\bar{s},e}(x)$, for some p -subsequence $\bar{s} = \langle i_1, \dots, i_p \rangle$ of $\langle 1, \dots, k \rangle$ and some $e \in \{0, 1\}^k$. We have to compute the limit of the probability

$$d_{2n}(\varphi_{\bar{s},e} : \text{ntr}) = \frac{|\mathbf{S}_{2n}(\varphi_{\bar{s},e} : \text{ntr})|}{|\mathbf{S}_{2n}(L)|}.$$

According to the assumption we made above, that we work only with 1-1 models of L , each L -structure \mathcal{M} is determined by a k -tuple $\langle W_1, \dots, W_k \rangle$ of elements of $\mathcal{P}(M)$, rather than a k -element subset $\{W_1, \dots, W_k\}$. So the number of 1-1 models is the number of k -tuples of distinct elements of $\mathcal{P}(M)$, i.e., $(2^n)_k$, thus, by (6),

$$|\mathbf{S}_{2n}(L)| = (2^{2n})_k \sim 2^{2kn}. \quad (7)$$

In order to compute $|\mathbf{S}_{2n}(\varphi_{\bar{s},e} : \text{ntnr})|$ we fix temporarily a set $A \subseteq M$ such that $|A| = n$. Since $\varphi_{\bar{s},e}(\mathcal{M}) = W_{i_1}^{e(i_1)} \cap \dots \cap W_{i_p}^{e(i_p)}$, we set $Z(A) = \left\{ \langle W_1, \dots, W_k \rangle : W_{i_1}^{e(i_1)} \cap \dots \cap W_{i_p}^{e(i_p)} = A \right\}$. Then clearly

$$|\mathbf{S}_{2n}(\varphi_{\bar{s},e} : \text{ntnr})| = |Z(A)| \cdot \binom{2n}{n}. \quad (8)$$

Now, $|Z(A)| = |Z_1(A)| \cdot |Z_2(A)|$, where

$$\begin{aligned} Z_1(A) &= \left\{ \langle W_{i_1}^{e(i_1)}, \dots, W_{i_p}^{e(i_p)} \rangle : W_{i_1}^{e(i_1)} \cap \dots \cap W_{i_p}^{e(i_p)} = A \right\} \quad \text{and} \\ Z_2(A) &= \left\{ \langle W_{j_1}^{e(j_1)}, \dots, W_{j_{k-p}}^{e(j_{k-p})} \rangle : \left\{ W_{j_1}^{e(j_1)}, \dots, W_{j_{k-p}}^{e(j_{k-p})} \right\} \subseteq \mathcal{P}(M) \setminus \left\{ W_{i_1}^{e(i_1)}, \dots, W_{i_p}^{e(i_p)} \right\} \right\}. \end{aligned}$$

In order to compute (or find upper bounds for) $|Z_1(A)|$ and $|Z_2(A)|$, we must distinguish the cases $p = 1$ and $2 \leq p \leq k$.

Case 1. $p = 1$. Without loss of generality we may assume that \bar{s} is the 1-subsequence (1), i.e., $\varphi_{\bar{s},e}(x)$ is either $U_1(x)$ or $-U_1(x)$. By Fact 7 (i), it suffices to consider only $U_1(x)$. Then $U_1(\mathcal{M}) = W_1$, so $Z_1(A) = \{W_1 : W_1 = A\}$, and hence $|Z_1(A)| = 1$. Also $Z_2(A) = \{ \langle W_2, \dots, W_k \rangle : \{W_2, \dots, W_k\} \subseteq \mathcal{P}(M) \setminus \{W_1\} \}$, so, in view of (6), $|Z_2(A)| = (2^{2n} - 1)_{k-1} \sim 2^{2(k-1)n}$. Thus, also $|Z(A)| = |Z_2(A)| \sim 2^{2(k-1)n}$, and therefore, letting A range over all sets of cardinality n , $\mathbf{S}_{2n}(U_1(x) : \text{ntnr}) \sim \binom{2n}{n} \cdot 2^{2(k-1)n}$. From the last equality, (7) and Fact 9 we get

$$d_{2n}(U_1(x) : \text{ntnr}) \sim \binom{2n}{n} \cdot \frac{2^{2(k-1)n}}{2^{kn}} \sim \binom{2n}{n} \cdot \frac{1}{2^{2n}} \leq \frac{4^n}{\sqrt{\pi n}} \cdot \frac{1}{2^{2n}} = \frac{1}{\sqrt{\pi n}} \rightarrow_n 0.$$

Therefore, $U_1(x)$ is regular.

Case 2. $2 \leq p \leq k$. Fix a p -subsequence of $\langle 1, \dots, k \rangle$ and an $e \in \{0, 1\}^p$. Then $\varphi_{\bar{s},e}(\mathcal{M}) = W_{i_1}^{e(i_1)} \cap \dots \cap W_{i_p}^{e(i_p)}$. Fixing temporarily a set $A \subseteq M$, as before $Z_2(A)$ consists of the $(k-p)$ -tuples of $\mathcal{P}(M) \setminus \left\{ W_{i_1}^{e(i_1)}, \dots, W_{i_p}^{e(i_p)} \right\}$, so $|Z_2(A)| = (2^{2n} - p)_{k-p} \sim 2^{2(k-p)n}$. The main difference of this case from the previous one lies in the computation of $Z_1(A)$. Observe that $W_{i_1}^{e(i_1)} \cap \dots \cap W_{i_p}^{e(i_p)} = A$ implies $A \subseteq W_{i_j}^{e(i_j)}$, for each $j = 1, \dots, p$. For each i_j we consider the cases $e(i_j) = 1$ and $e(i_j) = 0$.

- (i) If $e(i_j) = 1$, then $A \subseteq W_{i_j}$, so $W_{i_j} = A \cup Y_j$, for some $Y_j \subseteq M \setminus A$.
- (ii) If $e(i_j) = 0$, then $A \subseteq M \setminus W_{i_j}$, so $M \setminus W_{i_j} = A \cup Y_j$ for some $Y_j \subseteq M \setminus A$.

Thus, in both cases for each $j = 1, \dots, p$ there are at most as many possible choices for W_{i_j} as are the choices for $Y_j \subseteq M \setminus A$, i.e., $2^{|M \setminus A|} = 2^n$. In fact the choices of such Y_{i_j} 's are not independent. They must satisfy the condition $\bigcap_{j=1}^p Y_{i_j} = \emptyset$ (since otherwise $\bigcap_{j=1}^p W_{i_j}^{e(i_j)} \neq A$). Nevertheless, the number of possible p -tuples of $Z_1(A)$ are at most as many as the p -tuples of $\mathcal{P}(M \setminus A)$, i.e., $(2^n)_p \sim 2^{pn}$. Therefore, $|Z_1(A)| \leq (2^n)_p \sim 2^{pn}$. So $|Z(A)| = |Z_1(A)| \cdot |Z_2(A)| \lesssim 2^{2(k-p)n} \cdot 2^{pn} \sim 2^{2(k-p)n}$. Letting A range over all sets of cardinality n , we have

$$|\mathbf{S}_{2n}(\varphi_{\bar{s},e} : \text{ntnr})| \lesssim 2^{2(k-p)n} \cdot \binom{2n}{n}.$$

So, by (7) and Fact 9,

$$d_{2n}(\varphi_{\bar{s},e} : \text{ntnr}) \lesssim \frac{2^{2(k-p)n} \cdot \binom{2n}{n}}{2^{2kn}} \sim \frac{\binom{2n}{n}}{2^{pn}} \leq \frac{4^n}{2^{pn} \cdot \sqrt{\pi n}}.$$

Since $p \geq 2$, $\frac{4^n}{2^{pn} \cdot \sqrt{\pi n}} \leq \frac{4^n}{4^n \cdot \sqrt{\pi n}} = \frac{1}{\sqrt{\pi n}}$. So $d_{2n}(\varphi_{\bar{s},e} : \text{ntnr}) \rightarrow_n 0$, according to Fact 10. This completes the proof. \square

It is still open however whether *every* property of L , i.e., every $\varphi_E(x) = \bigvee_{e \in E} \varphi_e(x)$, for $E \subseteq \{0, 1\}^k$, is regular.

Question 13. Is every property $\varphi_E(x)$ of $L = \{U_1, \dots, U_k\}$ regular?

More generally:

Question 14. Is every property of a finite relational language regular?

3.2 A necessary condition for regularity

Next we shall give a necessary condition in order for a property $\varphi(x)$, (a) to have typicality degree 0, and (b) to be regular.

Let L be a language not necessarily relational. For any L -property $\varphi(x)$ and every $m \geq 1$ let us set

$$\varphi^{(m)} := (\exists x_1 \cdots \exists x_m) \left(\left(\bigwedge_{i \neq j} x_i \neq x_j \right) \wedge \left(\bigwedge_{i=1}^m \varphi(x_i) \right) \right).$$

The formula $\varphi^{(m)}$ is a sentence and says that $\varphi(x)$ is satisfied by at least m objects. So $m < k$ implies $\varphi^{(k)} \rightarrow \varphi^{(m)}$, and thus for every $m < k \leq n$,

$$\text{Mod}_n(\varphi^{(k)}) \subseteq \text{Mod}_n(\varphi^{(m)}). \quad (9)$$

As usual for every n let $\lfloor \frac{n}{2} \rfloor$ be the greatest integer $\leq n/2$. Then notice that by the definition of $\mathbf{S}_n(\varphi : \text{typ})$, for every $\mathcal{M} \in \mathbf{S}_n(L)$, $\mathcal{M} \in \mathbf{S}_n(\varphi : \text{typ})$ iff $\mathcal{M} \models \varphi^{(\lfloor \frac{n}{2} \rfloor + 1)}$, or

$$\mathbf{S}_n(\varphi : \text{typ}) = \text{Mod}_n(\varphi^{(\lfloor \frac{n}{2} \rfloor + 1)}). \quad (10)$$

On the other hand, for every \mathcal{M} and $k \leq |\mathcal{M}|$, $|\varphi(\mathcal{M})| = k$ implies $\mathcal{M} \models \varphi^{(k)}$, so $\mathcal{M} \in \mathbf{S}_{2n}(\varphi : \text{nt})$ implies $\mathcal{M} \models \varphi^{(n)}$, or

$$\mathbf{S}_{2n}(\varphi : \text{nt}) \subseteq \text{Mod}_{2n}(\varphi^{(n)}). \quad (11)$$

Lemma 15. Let L be any language and $\varphi(x)$ be an L -property. If there is $m \geq 1$ such that $\mu(\varphi^{(m)}) = 0$, then:

- (i) $d(\varphi : \text{typ}) = 0$, and
- (ii) $\varphi(x)$ is regular.

Proof. Let $\mu(\varphi^{(m)}) = 0$ for some fixed m . It means that

$$\mu_n(\varphi^{(m)}) = \frac{|\text{Mod}_n(\varphi^{(m)})|}{|\mathbf{S}_n(L)|} \rightarrow_n 0. \quad (12)$$

(i) For all $n \geq 2m$ we have $m < \lfloor \frac{n}{2} \rfloor + 1$, so by (9) and (10), $\mathbf{S}_n(\varphi : \text{typ}) = \text{Mod}_n(\varphi^{(\lfloor \frac{n}{2} \rfloor + 1)}) \subseteq \text{Mod}_n(\varphi^{(m)})$. Consequently, for every $n \geq 2m$,

$$\frac{|\mathbf{S}_n(\varphi : \text{typ})|}{|\mathbf{S}_n(L)|} \leq \frac{|\text{Mod}_n(\varphi^{(m)})|}{|\mathbf{S}_n(L)|}. \quad (13)$$

Then (13) combined with (12) yields

$$\frac{|\mathbf{S}_n(\varphi : \text{typ})|}{|\mathbf{S}_n(L)|} \rightarrow_n 0,$$

i.e., $d(\varphi : \text{typ}) = 0$.

(ii) For every $n \geq m$, by (9), (11) and (12) we have

$$d_{2n}(\varphi : \text{ntr}) = \frac{|\mathbf{S}_{2n}(\varphi : \text{ntr})|}{|\mathbf{S}_{2n}(L)|} \leq \frac{|\text{Mod}_{2n}(\varphi^{(n)})|}{|\mathbf{S}_{2n}(L)|} \leq \frac{|\text{Mod}_{2n}(\varphi^{(m)})|}{|\mathbf{S}_{2n}(L)|} \xrightarrow{n} 0.$$

Thus, $\varphi(x)$ is regular. \square

Corollary 16. If $d(\varphi : \text{typ}) = 1$, then $(\forall m \geq 1)(\mu(\varphi^{(m)}) = 1)$.

Proof. Assume $d(\varphi : \text{typ}) = 1$. First note that if L is relational, then the claim follows immediately from 15 using Theorem 2 about the 0-1 law for sentences of a relational L . For $d(\varphi : \text{typ}) = 1$ implies $d(\varphi : \text{typ}) \neq 0$, so, by Lemma 15, $(\forall m \geq 1)(\mu(\varphi^{(m)}) \neq 0)$ is true, and hence, by Theorem 2, $(\forall m \geq 1)(\mu(\varphi^{(m)}) = 1)$.

However the claim can be shown without appeal to Theorem 2, by a direct argument similar to that of Lemma 15. Namely, assuming $d(\varphi : \text{typ}) = 1$ we have

$$\frac{|\mathbf{S}_n(\varphi : \text{typ})|}{|\mathbf{S}_n(L)|} \xrightarrow{n} 1, \tag{14}$$

and since by (13) above, we have that for every $n \geq 2m$,

$$\frac{|\mathbf{S}_n(\varphi : \text{typ})|}{|\mathbf{S}_n(L)|} \leq \frac{|\text{Mod}_n(\varphi^{(m)})|}{|\mathbf{S}_n(L)|},$$

it follows that for all $m \geq 1$,

$$\frac{|\text{Mod}_n(\varphi^{(m)})|}{|\mathbf{S}_n(L)|} \xrightarrow{n} 1,$$

i.e., $(\forall m \geq 1)(\mu(\varphi^{(m)}) = 1)$. \square

4 The 0-1 law for typicality degrees and its failure for languages with unary properties

Given a language L , the 0-1 law for typicality degrees of properties of L can be defined by complete analogy with the corresponding law for sentences described in Section 2.1, as follows.

Definition 17. Let L be a finite language. We say that *the 0-1 law holds for typicality degrees of properties of L* , if for every L -property $\varphi(x)$, either $d(\varphi : \text{typ}) = 0$ or $d(\varphi : \text{typ}) = 1$.

We show in this section that the 0-1 law fails for the language $L = \{U_1, \dots, U_k\}$ with k unary predicates. Recall from the previous section the definition of a basic property $\varphi_{\bar{s},e}(x)$ of L , for a p -subsequence \bar{s} of $\langle 1, \dots, k \rangle$ and an $e \in \{0, 1\}^k$.

Theorem 18. Let $L = \{U_1, \dots, U_k\}$ with $k \geq 1$, and $\varphi_{\bar{s},e}(x)$ be a basic property of L .

- (i) If \bar{s} is a 1-sequence, then $d(\varphi_{\bar{s},e} : \text{typ}) = 1/2$.
- (ii) If \bar{s} is a p -sequence for $p \geq 2$, then $d(\varphi_{\bar{s},e} : \text{typ}) = 0$.

The subcase for $p = 2$ of this theorem requires special treatment, and it is more convenient to consider it separately. So we shall split Theorem 18 into Lemmas 19 and 20 below. The proof of the theorem is an immediate consequence of these lemmas.

Lemma 19.

- (i) If \bar{s} is a 1-sequence, then $d(\varphi_{\bar{s},e} : \text{typ}) = 1/2$.
- (ii) If \bar{s} is a p -sequence for $p \geq 3$, then $d(\varphi_{\bar{s},e} : \text{typ}) = 0$.

Proof. Let $\mathcal{M} = \langle M, W_1, \dots, W_k \rangle$ be an L -structure with $|M| = n$, and fix a property $\varphi_{\bar{s},e}(x)$ as above. Recall that given $\varphi_{\bar{s},e}(\mathcal{M}) = W_{i_1}^{e(i_1)} \cap \dots \cap W_{i_p}^{e(i_p)}$. The proof has many similarities with the proof of Theorem 12. Given any set $A \subseteq M$ let us define $Z(A)$, $Z_1(A)$, $Z_2(A)$ exactly as in the aforementioned proof. Then, as before, $|Z(A)| = |Z_1(A)| \cdot |Z_2(A)|$.

Setting $Z(m) = \bigcup \{Z(A) : |A| = m\}$, we have $|Z(m)| = \binom{n}{m} \cdot |Z(A)|$, for any A with $|A| = m$, and also

$$|\mathbf{S}_n(\varphi_{\bar{s},e} : \text{typ})| = \sum_{m > n/2} |Z(m)|. \quad (15)$$

(i) Let \bar{s} be a 1-sequence. Without loss of generality we assume that $\varphi_{\bar{s},e}(x) = U_1(x)$, so $\varphi_{\bar{s},e}(\mathcal{M}) = W_1$. Arguing exactly as in the proof of Theorem 12, we see that $|Z_1(A)| = 1$, and $|Z_2(A)| = (2^n - 1)_{k-1} \sim 2^{(k-1)n}$. Thus, also $|Z(A)| = |Z_2(A)| \sim 2^{(k-1)n}$, and therefore

$$\mathbf{S}_n(\varphi_{\bar{s},e}(x) : \text{typ}) \sim \sum_{m > n/2} \binom{n}{m} \cdot 2^{(k-1)n} \sim 2^{n-1} \cdot 2^{(k-1)n},$$

hence by (7),

$$d_n(\varphi_{\bar{s},e} : \text{typ}) \sim \frac{2^{n-1} \cdot 2^{(k-1)n}}{2^{kn}} \sim \frac{2^{n-1}}{2^n} = \frac{1}{2}.$$

Thus, $d(\varphi_{\bar{s},e} : \text{typ}) = \frac{1}{2}$, according to Fact 10.

(ii) Let \bar{s} be a p -sequence with $3 \leq p \leq k$. Arguing as in the corresponding part of the proof of Theorem 12, for each A with $|A| = m$, the p -tuples of $Z_1(A)$ are as many as the p -tuples $\langle Y_{j_1}, \dots, Y_{j_p} \rangle$ of elements of $\mathcal{P}(M \setminus A)$ whose members must have empty intersections. The number of all such sequences is $(2^{n-m})_p$ and constitutes an upper bound of $|Z_1(A)|$, that is, $|Z_1(A)| \leq (2^{n-m})_p$. So $|Z_1(A)| \leq (2^{n-m})_p$, while $|Z_2(A)| = (2^n - p)_{k-p}$. Therefore, $|Z(A)| = |Z_1(A)| \cdot |Z_2(A)| \leq (2^n - p)_{k-p} \cdot (2^{n-m})_p$. Setting $Z(m) = \bigcup \{Z(A) : A \subseteq M \text{ \& } |A| = m\}$, we have $|Z(m)| \leq \binom{n}{m} \cdot (2^n - p)_{k-p} \cdot (2^{n-m})_p$. So

$$\begin{aligned} \mathbf{S}_n(\varphi_{\bar{s},e} : \text{typ}) &= \sum_{m > n/2} |Z(m)| \leq \sum_{m > n/2} \binom{n}{m} [(2^n - p)_{k-p} \cdot (2^{n-m})_p] \\ &= (2^n - p)_{k-p} \cdot \sum_{m > n/2} \binom{n}{m} (2^{n-m})_p \sim 2^{(k-p)n} \cdot \sum_{m > n/2} \binom{n}{m} (2^{n-m})_p. \end{aligned}$$

Since $|\mathbf{S}_n(L)| = (2^n)_k \sim 2^{kn}$, it follows from the last relation that

$$d_n(\varphi_{\bar{s},e} : \text{typ}) \lesssim \frac{1}{2^{pn}} \cdot \sum_{m > n/2} \binom{n}{m} (2^{n-m})_p.$$

Setting $n - m = i$, this is written

$$d_n(\varphi_{\bar{s},e} : \text{typ}) \lesssim \frac{1}{2^{pn}} \cdot \sum_{i < n/2} \binom{n}{i} (2^i)_p. \quad (16)$$

Note that for $p > 2^i$, i.e., for $i < \log_2 p$, $(2^i)_p = 0$, so the above sum is equal to $\sum_{\log_2 p \leq i < n/2} \binom{n}{i} (2^i)_p$, however for notational simplicity we let i range over all $i < n/2$.

Now, for every n , k , besides the relation $(n)_k \sim n^k$, the relation $(n)_k \leq n^k$ holds too. In particular $(2^i)_p \leq 2^{pi}$, and also for $i < n/2$, $2^{pi} < 2^{\frac{pn}{2}}$, so (16) implies

$$d_n(\varphi_{\bar{s},e} : \text{typ}) \lesssim \frac{2^{\frac{pn}{2}}}{2^{pn}} \cdot \sum_{i < n/2} \binom{n}{i} = \frac{1}{2^{\frac{pn}{2}}} \sum_{i < n/2} \binom{n}{i}.$$

Now, by Fact 11, $\sum_{i < n/2} \binom{n}{i} \sim 2^{n-1}$, therefore

$$d_n(\varphi_{\bar{s},e} : \text{typ}) \lesssim \frac{2^{n-1}}{2^{\frac{pn}{2}}}. \quad (17)$$

Since $p \geq 3$, we have

$$\frac{2^{n-1}}{2^{\frac{pn}{2}}} \leq \frac{2^{n-1}}{2^{\frac{3n}{2}}} = \frac{1}{2^{\frac{n}{2}+1}} \rightarrow_n 0,$$

so, by Fact 10, (17) yields $d_n(\varphi_{\bar{s},e} : \text{typ}) \rightarrow_n 0$. This proves clause (ii) and completes the proof of the lemma. \square

In the proof of the next lemma we shall make use of Stirling numbers of the second kind. Recall that they are denoted $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$, or $S(n, k)$, where, for $1 \leq k \leq n$, $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$ counts the number of partitions of $\{1, \dots, n\}$ into k nonempty parts. The explicit formula for $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$ is (see [5, p. 231] or [6])

$$\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} = \frac{1}{k!} \sum_{i=0}^k (-1)^{k-i} \binom{k}{i} i^n. \quad (18)$$

Lemma 20. If \bar{s} is a 2-sequence then $d(\varphi_{\bar{s},e} : \text{typ}) = 0$.

Proof. Fix a 2-subsequence $\bar{s} = \langle i_1, i_2 \rangle$ of $\langle 1, \dots, k \rangle$ and an $e \in \{0, 1\}^k$. For any set $A \subseteq M$,

$$Z_1(A) = \left\{ \left\langle W_{i_1}^{e(i_1)}, W_{i_2}^{e(i_2)} \right\rangle : W_{i_1}^{e(i_1)} \cap W_{i_2}^{e(i_2)} = A \right\},$$

so $W_{i_1}^{e(i_1)} = A \cup Y_1$ and $W_{i_2}^{e(i_2)} = A \cup Y_2$, where $Y_1, Y_2 \subseteq M \setminus A$ and $Y_1 \cap Y_2 = \emptyset$. It follows that $|Z_1(A)| = |Q(A)|$, where $Q(A) = \{ \langle Y_1, Y_2 \rangle : Y_1, Y_2 \subseteq M \setminus A \wedge Y_1 \cap Y_2 = \emptyset \}$. The pairs $\langle Y_1, Y_2 \rangle$ are of two kinds: those for which Y_1, Y_2 form a partition of $M \setminus A$, i.e., $Y_1 \cup Y_2 = M \setminus A$, and those for which $Y_1 \cup Y_2 \neq M \setminus A$. That is, $Q(A) = Q_1(A) \cup Q_2(A)$, where

$$\begin{aligned} Q_1(A) &= \{ \langle Y_1, Y_2 \rangle : Y_1 \cap Y_2 = \emptyset \wedge Y_1 \cup Y_2 = M \setminus A \}, \\ Q_2(A) &= \{ \langle Y_1, Y_2 \rangle : Y_1 \cap Y_2 = \emptyset \wedge Y_1 \cup Y_2 \neq M \setminus A \}, \\ |Z_1(A)| &= |Q(A)| = |Q_1(A)| + |Q_2(A)|. \end{aligned} \quad (19)$$

Now, $|Q_1(A)|$ and $|Q_2(A)|$ can be easily calculated in terms of the 2-partitions and 3-partitions of $M \setminus A$, respectively. Let $\Pi(M \setminus A, 2)$, $\Pi(M \setminus A, 3)$ denote the sets of partitions of $M \setminus A$ into 2 and 3 nonempty parts, respectively. Then $|\Pi(M \setminus A, 2)| = \left\{ \begin{smallmatrix} n-m \\ 2 \end{smallmatrix} \right\}$ and $|\Pi(M \setminus A, 3)| = \left\{ \begin{smallmatrix} n-m \\ 3 \end{smallmatrix} \right\}$. Now, a member of $\Pi(M \setminus A, 2)$ is a 2-element subset of $M \setminus A$, while $Q_1(A)$ consists of *ordered pairs* of such subsets, therefore $|Q_1(A)| = 2 \cdot |\Pi(M \setminus A, 2)| = 2 \cdot \left\{ \begin{smallmatrix} n-m \\ 2 \end{smallmatrix} \right\}$. Analogously every member of $\Pi(M \setminus A, 3)$ is a 3-element subset of $M \setminus A$, and each such subset provides $(3)_2$ pairs that belong to $Q_2(A)$, so $|Q_2(A)| = (3)_2 \cdot |\Pi(M \setminus A, 3)| = 6 \cdot \left\{ \begin{smallmatrix} n-m \\ 3 \end{smallmatrix} \right\}$. For every $n \geq 2$, it is easy to see (without appealing to (18)) that $\left\{ \begin{smallmatrix} n \\ 2 \end{smallmatrix} \right\} = 2^{n-1} - 1$, so for every A with $|A| = m \leq n - 2$, $|Q_1(A)| = 2 \cdot (2^{n-m-1} - 1) = 2^{n-m} - 2$. If $Q_1(m) = \bigcup \{ Q_1(A) : |A| = m \}$, then for $m \leq n - 2$,

$$|Q_1(m)| = \binom{n}{m} \cdot (2^{n-m} - 2). \quad (20)$$

On the other hand, for every $n \geq 3$, $\left\{ \begin{smallmatrix} n \\ 3 \end{smallmatrix} \right\}$ is calculated using formula (18), which yields $\left\{ \begin{smallmatrix} n \\ 3 \end{smallmatrix} \right\} = \frac{1}{6} [3 - 3 \cdot 2^n + 3^n]$. Therefore, for $|A| = m \leq n - 3$, $|Q_2(A)| = 3 - 3 \cdot 2^{n-m} + 3^{n-m}$. So setting as before $Q_2(m) = \bigcup \{ Q_2(A) : |A| = m \}$, we have for $m \leq n - 3$,

$$|Q_2(m)| = \binom{n}{m} \cdot (3 - 3 \cdot 2^{n-m} + 3^{n-m}). \quad (21)$$

Finally by (19), (20) and (21) above we obtain, for $|A| = m \leq n - 3$, $|Z_1(m)| = |Q_1(m)| + |Q_2(m)| = \binom{n}{m}(3^{n-m} - 2^{n-m+1} + 1)$. Recall also from above that for $p = 2$, $|Z_2(m)| \sim \binom{n}{m}2^{(k-2)n}$, so for $m \leq n - 3$,

$$|Z(m)| = |Z_1(m)| \cdot |Z_2(m)| \sim \binom{n}{m} \cdot [2^{(k-2)n} \cdot (3^{n-m} - 2^{n-m+1} + 1)]. \quad (22)$$

Now, it is easy to see that

$$\mathbf{S}_n(\varphi_{\bar{s},e} : \text{typ}) = \sum_{n \geq m > n/2} |Z(m)| \sim \sum_{n-3 \geq m > n/2} |Z(m)|,$$

so by (22),

$$\mathbf{S}_n(\varphi_{\bar{s},e} : \text{typ}) \sim 2^{(k-2)n} \cdot \sum_{n-3 \geq m > n/2} \binom{n}{m} \cdot (3^{n-m} - 2^{n-m+1} + 1),$$

and, given that $\mathbf{S}_n(L) = 2^{kn}$,

$$d_n(\varphi_{\bar{s},e} : \text{typ}) \sim \frac{1}{2^{2n}} \cdot \sum_{n-3 \geq m > n/2} \binom{n}{m} \cdot (3^{n-m} - 2^{n-m+1} + 1).$$

Setting $n - m = i$, this is written

$$\begin{aligned} d_n(\varphi_{\bar{s},e} : \text{typ}) &\sim \frac{1}{2^{2n}} \cdot \sum_{3 \leq i < n/2} \binom{n}{i} \cdot (3^i - 2^{i+1} + 1) \leq \frac{1}{2^{2n}} \cdot \sum_{3 \leq i < n/2} \binom{n}{i} \cdot (3^{\frac{n}{2}} - 2^{\frac{n}{2}+1} + 1) \\ &= \frac{3^{\frac{n}{2}} - 2^{\frac{n}{2}+1} + 1}{2^{2n}} \cdot \sum_{3 \leq i < n/2} \binom{n}{i} \leq \frac{3^{\frac{n}{2}} - 2^{\frac{n}{2}+1} + 1}{2^{2n}} \cdot \sum_{0 \leq i < n/2} \binom{n}{i}. \end{aligned}$$

By Fact 11, the last quantity is

$$\sim \frac{3^{\frac{n}{2}} - 2^{\frac{n}{2}+1} + 1}{2^{2n}} \cdot 2^{n-1} \sim \frac{3^{\frac{n}{2}} - 2^{\frac{n}{2}+1} + 1}{2^{n+1}} \sim \frac{1}{2} \left(\frac{3^{\frac{n}{2}}}{4^{\frac{n}{2}}} - \frac{1}{2^{\frac{n}{2}-1}} + \frac{1}{2^n} \right).$$

So finally,

$$d_n(\varphi_{\bar{s},e} : \text{typ}) \lesssim \frac{1}{2} \left(\frac{3^{\frac{n}{2}}}{4^{\frac{n}{2}}} - \frac{1}{2^{\frac{n}{2}-1}} + \frac{1}{2^n} \right) \rightarrow_n 0.$$

This completes the proof. \square

An immediate consequence of Theorem 18 (i) is the following.

Corollary 21. The 0-1 law for typicality degrees of properties of a relational language fails in general.

Let us consider at this point another question related to property “ $U(x)$ ”, namely the question about the probability of the sentences $U(x)^{(m)}$, for $m \geq 1$. We can see that, in contrast to the typicality degree $1/2$ of $U(x)$, the truth probability of the sentences $U(x)^{(m)}$ is 1.

Proposition 22. For every $m \geq 1$, $\mu(U(x)^{(m)}) = 1$.

Proof. The sentence $U(x)^{(m)}$ says that “ $U(x)$ is satisfied by at least m elements”. Therefore, $\text{Mod}_n(U(x)^{(m)}) = \{A \subseteq \{1, \dots, n\} : |A| \geq m\}$. Then $|\text{Mod}_n(U(x)^{(m)})| = 2^n - \sum_{0 \leq i < m} \binom{n}{i}$, and therefore

$$\mu_n(U(x)^{(m)}) = \frac{2^n - \sum_{0 \leq i < m} \binom{n}{i}}{2^n} = 1 - \frac{\sum_{0 \leq i < m} \binom{n}{i}}{2^n}.$$

The nominator $\sum_{0 \leq i < m} \binom{n}{i}$ of the fraction on the right-hand side is a polynomial in n of degree $m - 1$, so its quotient by the exponential 2^n goes to 0 as n grows. Thus, $\mu(U(x)^{(m)}) = \lim_{n \rightarrow \infty} \mu_n(U(x)^{(m)}) = 1$. \square

Question 23. Does the 0-1 law about typicality degrees hold for every property of a relational language without unary predicates?

We close this section with a remark about Question 23. Some people believe that the answer to this question must be positive on the basis that the rather general method of *extension axioms* that was used by R. Fagin [3] to prove the 0-1 law for truth degrees of *sentences* of a relational language, could be also applied somehow to the case of typicality degrees of properties.¹ The problem, however, is that this specific method works for sentences of *every* relational language, including those that contain unary predicates, while, as we saw above, in the case of typicality degrees the method should not work when unary predicates are included. We don't know how this gap could be bridged and if Fagin's method is actually applicable to the present case.

5 Languages with a single unary function symbol

The failure of 0-1 law for typicality degrees of properties of languages with unary predicates is a divergence from the behavior of truth probabilities of sentences. In this section we consider some properties of the language $L = \{F\}$ where F is a unary function symbol. We saw in Example 3 that $\mu(\forall x(F(x) \neq x)) = e^{-1}$, which means that the 0-1 law does not hold for sentences of L . The question is whether the 0-1 law fails also for typicality degrees of properties of this language. In this section we consider the property $\varphi(x) := (F(x) \neq x)$. It is shown that $\varphi(x)$ is regular and that the degree of $\varphi(x)$ is 1.

Proposition 24. Let $L = \{F\}$ and let $\varphi(x) := (F(x) \neq x)$. Then $d(\varphi : \text{typ}) = 1$.

Proof. Let $\mathcal{M} = \langle M, f \rangle$ with $|M| = n$ and $A \subseteq M$ such that $A = \varphi(\mathcal{M})$. Then $a \in A \Leftrightarrow f(a) \neq a$. Thus, if $G(A) = \{f \in M^M : \varphi(\mathcal{M}) = A\}$, every $f \in G(A)$ can be identified with the pair $(f \upharpoonright A, \text{id}_{(M \setminus A)})$, or since $\text{id}_{(M \setminus A)}$ is unique, with $f \upharpoonright (M \setminus A)$. As we argued in Example 3, if $|A| = m$ then $|G(A)| = (n-1)^m$. Let us set $G(m) = \bigcup \{G(A) : A \subseteq M \ \& \ |A| = m\}$. Then, since there are $\binom{n}{m}$ sets of cardinality m , $|G(m)| = \binom{n}{m}(n-1)^m$. Also by definition, $\mathbf{S}_n(\varphi : \text{typ}) = \bigcup \{G(m) : m > n/2\}$, therefore

$$|\mathbf{S}_n(\varphi : \text{typ})| = \sum_{m > n/2} |G(m)| = \sum_{m > n/2} \binom{n}{m} (n-1)^m. \quad (23)$$

It is more convenient to set $m = n - k$ and write this sum in the form

$$|\mathbf{S}_n(\varphi : \text{typ})| = \sum_{k < n/2} \binom{n}{n-k} (n-1)^{n-k} = \sum_{k < n/2} \binom{n}{k} (n-1)^{n-k},$$

or

$$|\mathbf{S}_n(\varphi : \text{typ})| = \sum_{k < n/2} a_k(n), \quad (24)$$

where $a_k(n) = \binom{n}{k} (n-1)^{n-k}$. Then

$$d_n(\varphi : \text{typ}) = \frac{|\mathbf{S}_n(\varphi : \text{typ})|}{n^n} = \sum_{k < n/2} \frac{a_k(n)}{n^n}.$$

Setting for simplicity, $a_n = \sum_{k < n/2} \frac{a_k(n)}{n^n}$, we have to compute the limit

$$d(\varphi : \text{typ}) = \lim_{n \rightarrow \infty} a_n. \quad (25)$$

¹ In contrast, the method of Glebskii et al. [4] seems to be rather ad hoc.

Now, $\frac{a_k(n)}{n^n} = \frac{1}{n^n} \cdot \binom{n}{k} (n-1)^{n-k}$, or, multiplying and dividing the right-hand side by $(n-1)^k$, this is written $\frac{a_k(n)}{n^n} = \frac{(n-1)^n}{n^n} \cdot \binom{n}{k} \frac{1}{(n-1)^k}$. Therefore,

$$a_n = \sum_{k < n/2} \left(\frac{n-1}{n} \right)^n \cdot \binom{n}{k} \frac{1}{(n-1)^k} = \left(\frac{n-1}{n} \right)^n \cdot \sum_{k < n/2} \binom{n}{k} \frac{1}{(n-1)^k} = b_n \cdot c_n,$$

where $b_n = \left(\frac{n-1}{n} \right)^n$ and $c_n = \sum_{k < n/2} \binom{n}{k} \frac{1}{(n-1)^k}$, and

$$\lim_{n \rightarrow \infty} a_n = \left(\lim_{n \rightarrow \infty} b_n \right) \cdot \left(\lim_{n \rightarrow \infty} c_n \right). \quad (26)$$

Now, $\lim_{n \rightarrow \infty} b_n = e^{-1}$. So it remains to compute

$$\lim_{n \rightarrow \infty} c_n = \lim_{n \rightarrow \infty} \sum_{k < n/2} \binom{n}{k} \frac{1}{(n-1)^k}. \quad (27)$$

We shall show that $\lim_{n \rightarrow \infty} c_n = e$. It suffices to show that $\lim_{n \rightarrow \infty} c_{2n} = \lim_{n \rightarrow \infty} c_{2n+1} = e$.

Claim 25. $\lim_{n \rightarrow \infty} c_{2n} = e$.

Proof of Claim. We have

$$c_{2n} = \sum_{k < n} \binom{2n}{k} \frac{1}{(2n-1)^k} = \sum_{k=0}^{n-1} \binom{2n}{k} \frac{1}{(2n-1)^k}.$$

However it is easy to see that $\lim_{n \rightarrow \infty} c_{2n} = \lim_{n \rightarrow \infty} c'_{2n}$, where

$$c'_{2n} = \sum_{k=0}^n \binom{2n}{k} \frac{1}{(2n-1)^k}.$$

This is because $c'_{2n} - c_{2n} = \binom{2n}{n} \frac{1}{(2n-1)^n}$, which goes to 0 when n goes to infinity, as follows easily from Fact 9. Therefore, it suffices to show that $\lim_{n \rightarrow \infty} c'_{2n} = e$. To show that, we compare each term c'_{2n} with the term

$$g_n = \sum_{k=0}^n \binom{n}{k} \frac{1}{(n-1)^k} = \left(1 + \frac{1}{n-1} \right)^n = \left(\frac{n}{n-1} \right)^n,$$

for which it is well-known that $\lim_{n \rightarrow \infty} g_n = e$. Specifically we compare each summand $A_k(n) = \binom{2n}{k} \frac{1}{(2n-1)^k}$ of c'_{2n} , for $k \leq n$, with the corresponding summand $B_k(n) = \binom{n}{k} \frac{1}{(n-1)^k}$ of g_n . Then we have

$$A_k(n) = \frac{1}{k!} \frac{(2n-k+1)(2n-k+2) \cdots 2n}{(2n-1)^k} = \frac{1}{k!} \frac{P(2n)}{Q(2n)},$$

while

$$B_k(n) = \frac{1}{k!} \frac{(n-k+1)(n-k+2) \cdots n}{(n-1)^k} = \frac{1}{k!} \frac{P(n)}{Q(n)},$$

where $P(x)$ and $Q(x)$ are polynomials of degree k . If α_k and β_k are the leading coefficients of $P(x)$ and $Q(x)$, respectively, then

$$\lim_{n \rightarrow \infty} \frac{P(2n)}{Q(2n)} = \lim_{n \rightarrow \infty} \frac{P(n)}{Q(n)} = \frac{\alpha_k}{\beta_k},$$

therefore $\lim_{n \rightarrow \infty} (A_k(n) - B_k(n)) = 0$. Besides,

$$c'_{2n} - g_n = \sum_{k=0}^n A_k(n) - \sum_{k=0}^n B_k(n) = \sum_{k=0}^n (A_k(n) - B_k(n)),$$

so

$$\lim_{n \rightarrow \infty} (c'_{2n} - g_n) = \sum_{k=0}^n \lim_{n \rightarrow \infty} (A_k(n) - B_k(n)) = 0.$$

Therefore, $\lim_{n \rightarrow \infty} c'_{2n} = \lim_{n \rightarrow \infty} g_n = e$, as required. \square

Claim 26. $\lim_{n \rightarrow \infty} c_{2n+1} = e$.

Proof of Claim. This is almost the same as in Claim 25. First notice that $k < (2n+1)/2 \Leftrightarrow k \leq n$, so

$$c_{2n+1} = \sum_{k < (2n+1)/2} \binom{2n+1}{k} \frac{1}{(2n)^k} = \sum_{k=0}^n \binom{2n+1}{k} \frac{1}{(2n)^k}.$$

Then we set for every $k \leq n$,

$$A'_k(n) = \binom{2n+1}{k} \frac{1}{(2n)^k} = \frac{1}{k!} \frac{P(2n+1)}{Q(2n+1)}$$

and compare it again with

$$B_k(n) = \frac{1}{k!} \frac{P(n)}{Q(n)}$$

of the preceding case. As before we have

$$\lim_{n \rightarrow \infty} \frac{P(2n+1)}{Q(2n+1)} = \lim_{n \rightarrow \infty} \frac{P(n)}{Q(n)} = \frac{\alpha_k}{\beta_k},$$

therefore $\lim_{n \rightarrow \infty} (A'_k(n) - B_k(n)) = 0$, and hence

$$\lim_{n \rightarrow \infty} (c_{2n+1} - g_n) = \sum_{k=0}^n \lim_{n \rightarrow \infty} (A'_k(n) - B_k(n)) = 0.$$

So $\lim_{n \rightarrow \infty} c_{2n+1} = \lim_{n \rightarrow \infty} g_n = e$. \square

This completes the proof of Proposition 24. \square

Proposition 27. The property $\varphi(x) := (F(x) \neq x)$ is regular.

Proof. Let $\mathcal{M} = \langle M, f \rangle$ be an L -structure with $|M| = 2n$. Then $\mathcal{M} \in \mathbf{S}_{2n}(\varphi : \text{ntnr})$ if and only if $|\varphi(\mathcal{M})| = n$. Let $A = \varphi(\mathcal{M})$. Then $A = \{a \in M : f(a) \neq a\}$ and hence $a \notin A \Leftrightarrow f(a) = a$, that is, $f \upharpoonright (M \setminus A) = \text{id}_{M \setminus A}$. Since $\text{id}_{M \setminus A}$ is unique, it follows that if $G(A)$ is the set of $f \in M^M$ such that $\varphi(\mathcal{M}) = A$, then

$$|G(A)| = |\{f \upharpoonright A : \varphi(\mathcal{M}) = A\}| = |\{g \in M^A : \forall x (g(x) \neq x)\}|.$$

Since $|M| = 2n$ and $|A| = n$ we have, as argued in Example 3, $|G(A)| = (2n-1)^n$. Since there are $\binom{2n}{n}$ such sets A , it follows that

$$|\mathbf{S}_{2n}(\varphi : \text{ntnr})| = \left| \bigcup \{G(A) : |A| = n\} \right| = \binom{2n}{n} \cdot (2n-1)^n.$$

On the other hand $|\mathbf{S}_{2n}(L)| = (2n)^{2n}$, hence

$$\frac{|\mathbf{S}_{2n}(\varphi : \text{ntnr})|}{|\mathbf{S}_{2n}(L)|} = \frac{\binom{2n}{n} \cdot (2n-1)^n}{(2n)^{2n}} \leq \frac{\binom{2n}{n} \cdot (2n)^n}{(2n)^{2n}} = \frac{\binom{2n}{n}}{(2n)^n}.$$

By Fact 9, $\binom{2n}{n} \leq \frac{4^n}{\sqrt{\pi n}}$, so the preceding inequality implies

$$\frac{|\mathbf{S}_{2n}(\varphi : \text{ntnr})|}{|\mathbf{S}_{2n}(L)|} \leq \frac{4^n}{(2n)^n \cdot \sqrt{\pi n}} = \frac{2^n}{n^n \cdot \sqrt{\pi n}} \rightarrow_n 0. \quad \square$$

Corollary 28. $d(F(x) = x : \text{typ}) = 0$.

Proof. By Proposition 27, $F(x) = x$ is regular, while by Proposition 24, $d(F(x) \neq x : \text{typ}) = 1$. Therefore, in view of Fact 7 (iii), $d(F(x) = x : \text{typ}) = 1 - d(F(x) \neq x : \text{typ}) = 0$. \square

The next proposition shows that the criterion of Lemma 15 (ii), which allows one to deduce the regularity of property $\varphi(x)$ whenever for some $m \geq 1$, $\mu(\varphi^{(m)}) = 0$, cannot be used for properties $F(x) \neq x$ and $F(x) = x$.

Proposition 29. For every $m \geq 1$,

- (i) $\mu((F(x) \neq x)^{(m)}) = 1$.
- (ii) $\frac{e^{-1}}{m!} \leq \mu((F(x) = x)^{(m)}) \leq \frac{1}{m!}$. Moreover, $\mu((F(x) = x)^{(1)}) = 1 - e^{-1}$.

Proof. (i) By definition $(F(x) \neq x)^{(m)}$ holds in $\langle M, f \rangle$ if and only if there are distinct $x_1, \dots, x_m \in M$ such that $f(x_i) \neq x_i$ for every $i = 1, \dots, m$. Equivalently, if and only if for every $A \subseteq M$ such that $f \upharpoonright A = \text{id}$, $|A| \leq |M| - m$. Fixing M with $|M| = n$, for a given $A \subseteq M$ the totality of f such that $f \upharpoonright A = \text{id}$ are $n^{n-|A|}$, while when A ranges over all subsets with $|A| \leq n - m$, the totality of f for which $\langle M, f \rangle$ satisfies $(F(x) \neq x)^{(m)}$ has cardinality

$$|\text{Mod}_n((F(x) \neq x)^{(m)})| = \sum_{|A| \leq n-m} n^{n-|A|} = \sum_{m \leq i \leq n} n^i.$$

Therefore,

$$\mu_n((F(x) \neq x)^{(m)}) = \frac{\sum_{m \leq i \leq n} n^i}{n^n} \xrightarrow[n]{} 1.$$

(ii) Let us show first the second claim, that $\mu((F(x) = x)^{(1)}) = 1 - e^{-1}$. This follows from Example 3, and the fact that for every sentence φ , $\mu(\neg\varphi) = 1 - \mu(\varphi)$. So

$$\begin{aligned} \mu((F(x) = x)^{(1)}) &= \mu(\exists x(F(x) = x)) = \mu(\neg(\forall x)(F(x) \neq x)) \\ &= 1 - \mu(\forall x(F(x) \neq x)) = 1 - \frac{1}{e} = \frac{e-1}{e}. \end{aligned}$$

Now, consider the sentence $(F(x) = x)^{(m)}$ for $m \geq 1$. The sentence $(F(x) = x)^{(m)}$ holds in $\langle M, f \rangle$ if and only if there are distinct $x_1, \dots, x_m \in M$ such that $f(x_i) = x_i$ for every $i = 1, \dots, m$, i.e., if there is $A \subseteq M$ with $|A| = m$ such that $f \upharpoonright A = \text{id}$. If $|M| = n$, for each A with $|A| = m$ there are n^{n-m} functions such that $f \upharpoonright A = \text{id}$. Since there are $\binom{n}{m}$ such sets A with $|A| = m$, the totality of functions of this kind is at most $\binom{n}{m} n^{n-m}$ (because this totality possibly contains repetitions). So $|\text{Mod}_n((F(x) = x)^{(m)})| \leq \binom{n}{m} n^{n-m}$.

On the other hand, for a fixed A with $|A| = m$, let X_A be the collection of functions f such that $f \upharpoonright A = \text{id}$ and $f(x) \neq x$ for every $x \in M \setminus A$. Since for every $f \in X_A$, $f(x)$ may take independently $n - 1$ possible values on $M \setminus A$, it follows that $|X_A| = (n - 1)^{n-m}$ and, moreover, if $A \neq A'$ then $X_A \cap X_{A'} = \emptyset$. So $|\text{Mod}_n((F(x) = x)^{(m)})| \geq \binom{n}{m} (n - 1)^{n-m}$. Therefore,

$$\frac{\binom{n}{m} (n - 1)^{n-m}}{n^n} \leq \mu_n((F(x) = x)^{(m)}) \leq \frac{\binom{n}{m} n^{n-m}}{n^n}. \quad (28)$$

Denoting by l_n the preceding lower bound in (28) we have

$$\begin{aligned} l_n &= \frac{\binom{n}{m} (n - 1)^{n-m}}{n^n} = \frac{\binom{n}{m}}{(n - 1)^m} \cdot \frac{(n - 1)^n}{n^n} \\ &= \frac{1}{m!} \frac{(n - m + 1)(n - m + 2) \cdots (n - 1)n}{(n - 1)^m} \cdot \left(\frac{n - 1}{n}\right)^n. \end{aligned}$$

Therefore,

$$\lim_{n \rightarrow \infty} l_n = \frac{1}{m!} \lim_{n \rightarrow \infty} \frac{(n-m+1)(n-m+2) \cdots (n-1)n}{(n-1)^m} \cdot \lim_{n \rightarrow \infty} \left(\frac{n-1}{n} \right)^n.$$

Now, $\lim_{n \rightarrow \infty} \frac{(n-m+1)(n-m+2) \cdots (n-1)n}{(n-1)^m} = 1$, because both the nominator and the denominator are polynomials of degree m with leading coefficients 1, while $\lim_{n \rightarrow \infty} \left(\frac{n-1}{n} \right)^n = e^{-1}$. Therefore, $\lim_{n \rightarrow \infty} l_n = \frac{e^{-1}}{m!}$.

Similarly, if $u_n = \frac{\binom{n}{m} n^{n-m}}{n^n}$ is the upper bound in (28) above, then

$$u_n = \frac{\binom{n}{m}}{n^m} = \frac{1}{m!} \cdot \frac{(n-m+1)(n-m+2) \cdots (n-1)n}{n^m},$$

and comparing as before the polynomials of the fraction we find $\lim_{n \rightarrow \infty} u_n = \frac{1}{m!}$. So finally

$$\frac{e^{-1}}{m!} \leq \mu((F(x) = x)^{(m)}) = \lim_{n \rightarrow \infty} \mu_n((F(x) = x)^{(m)}) \leq \frac{1}{m!}.$$

Notice that the value $1 - e^{-1} = \frac{e-1}{e}$ of $\mu((F(x) = x)^{(1)})$ conforms with the general bounds given above, since $e^{-1} \leq \frac{e-1}{e} \leq 1$. \square

Corollary 30. The converse of Lemma 15 (i) is false for the language $L = \{F\}$. Namely, there is $\varphi(x)$ of L such that $d(\varphi : \text{typ}) = 0$ while $\forall m \geq 1 \mu(\varphi^{(m)}) > 0$.

Proof. Take $\varphi(x) : (F(x) = x)$. By Corollary 28, $d(F(x) = x : \text{typ}) = 0$, while by Proposition 29, for every $m \geq 1 \mu((F(x) = x)^{(m)}) \geq \frac{e^{-1}}{m!}$. \square

6 Degrees of some properties of graphs

In this section we examine the typicality degree of some natural first-order properties of finite *undirected* graphs. We stress the fact that we deal with undirected graphs only, since they constitute a tiny minority among all graphs or, in more technical terms, they have density zero in the class of all graphs.

The language of graphs is $L = \{E\}$, where E is the symbol of a binary symmetric and irreflexive relation. Thus, an L -structure is a pair $G = \langle A, E \rangle$, where A is the set of nodes of G and E is its *adjacency* relation; xEy means that “the nodes x, y are adjacent”, i.e., connected with an edge. By assumption $xEy \Leftrightarrow yEx$ and for all $x \in A \neg(xEx)$. So E is a set of 2-element subsets of A , that is $E \subseteq [A]^2$, and $G \models xEy \Leftrightarrow \{x, y\} \in E$. So if $|A| = n$, then $|[A]^2| = \binom{n}{2}$, therefore $|\mathbf{S}_n(L)| = 2^{\binom{n}{2}}$.

Let us consider the following examples of first-order properties of L :

- (1) $\varphi_{\text{none}}(x)$: “ x is an isolated node” $[(\forall y)\neg(xEy)]$.
- (2) $\varphi_{\text{all}}(x)$: “ x is adjacent to every node” $[(\forall y)(xEy)]$.
- (3) $\varphi_{\text{one}}(x)$: “ x is adjacent to exactly one node” $[(\exists!y)(xEy)]$.

Proposition 31. Properties $\varphi_{\text{none}}(x)$, $\varphi_{\text{all}}(x)$ and $\varphi_{\text{one}}(x)$ have typicality degree 0. That is, $d(\varphi_{\text{none}} : \text{typ}) = d(\varphi_{\text{all}} : \text{typ}) = d(\varphi_{\text{one}} : \text{typ}) = 0$.

Proof. We shall prove the claim using Lemma 15 (i), namely it suffices to prove that $\mu(\varphi_{\text{none}}^{(1)}) = \mu(\varphi_{\text{all}}^{(1)}) = \mu(\varphi_{\text{one}}^{(1)}) = 0$.

We first prove that $\mu(\varphi_{\text{none}}^{(1)}) = 0$. We have $\varphi_{\text{none}}^{(1)} := (\exists x)(\forall y)\neg(xEy)$. We must show that

$$\frac{|\text{Mod}_n(\varphi_{\text{none}}^{(1)})|}{|\mathbf{S}_n(L)|} \rightarrow_n 0. \tag{29}$$




Figure 1: The graph G contains an isolated node a .

We saw above that $|\mathbf{S}_n(L)| = 2^{\binom{n}{2}}$, so we have to estimate also $|\text{Mod}_n(\varphi_{\text{none}}^{(1)})|$. Let $G = \langle A, E \rangle$ be a graph with n nodes satisfying $\varphi_{\text{none}}^{(1)}$. Then G contains at *least one* isolated node a (see Figure 1).

It follows that if $A' = A \setminus \{a\}$ and $G' = \langle A', E' \rangle$, where E' is the restriction of E to A' , then G' can be *any* graph with $n - 1$ nodes (including those with any number of isolated nodes). Therefore, there are $2^{\binom{n-1}{2}}$ different graphs on A in which a is an isolated node. Letting a range over every node of A , it follows that the total number of graphs in $\mathbf{S}_n(L)$ that satisfy $\varphi_{\text{none}}^{(1)}$, i.e., $|\text{Mod}_n(\varphi_{\text{none}}^{(1)})|$, is at most $n2^{\binom{n-1}{2}}$. Thus,

$$\frac{|\text{Mod}_n(\varphi_{\text{none}}^{(1)})|}{|\mathbf{S}_n(L)|} \leq \frac{n2^{\binom{n-1}{2}}}{2^{\binom{n}{2}}} = \frac{n2^{\frac{(n-2)(n-1)}{2}}}{2^{\frac{n(n-1)}{2}}} = \frac{n}{2^{n-1}} \rightarrow_n 0,$$

so (29) is true.

We will now prove that $\mu(\varphi_{\text{all}}^{(1)}) = 0$. We have $\varphi_{\text{all}}^{(1)} := (\exists x)(\forall y)(xEy)$ and we must show that

$$\frac{|\text{Mod}_n(\varphi_{\text{all}}^{(1)})|}{|\mathbf{S}_n(L)|} \rightarrow_n 0. \quad (30)$$

The argument is quite similar to that of the previous case. If $G = \langle A, E \rangle$ is a graph with n nodes satisfying $\varphi_{\text{all}}^{(1)}$, there is $a \in A$ which is adjacent to all other nodes of G (see Figure 2).




Figure 2: The graph G contains a node a adjacent to all other nodes of G .

It means that if $A' = A \setminus \{a\}$ and $G' = \langle A', E' \rangle$ is as before, then G' can be again of any possible form, so as before

$$\frac{|\text{Mod}_n(\varphi_{\text{all}}^{(1)})|}{|\mathbf{S}_n(L)|} \leq \frac{n}{2^{n-1}} \rightarrow_n 0.$$

Finally, we prove that $\mu(\varphi_{\text{one}}^{(1)}) = 0$. We have $\varphi_{\text{one}}^{(1)} := (\exists x)(\exists! y)(xEy)$ and we have to show that

$$\frac{|\text{Mod}_n(\varphi_{\text{one}}^{(1)})|}{|\mathbf{S}_n(L)|} \xrightarrow{n} 0. \quad (31)$$

Let $G = \langle A, E \rangle$ be a graph, with $|A| = n$, satisfying $\varphi_{\text{one}}^{(1)}$. Pick an $a \in A$ witnessing $\varphi_{\text{one}}^{(1)}$. If again $A' = A \setminus \{a\}$, a is adjacent to a unique element $b \in A'$ (see Figure 3), so there are $n - 1$ choices for the given a , to be connected.




Figure 3: The node a is adjacent to the unique element $b \in A' = A \setminus \{a\}$.

For each such choice the subgraph $G' = \langle A', E' \rangle$ can have any possible form, so the number of graphs on A in which the specific a witnesses $\varphi_{\text{one}}^{(1)}$ is at most $(n - 1) \cdot 2^{\binom{n-1}{2}}$. Letting again a range over A , it follows that $|\text{Mod}_n(\varphi_{\text{one}}^{(1)})| \leq n(n - 1)2^{\binom{n-1}{2}}$. So

$$\frac{|\text{Mod}_n(\varphi_{\text{one}}^{(1)})|}{|\mathbf{S}_n(L)|} \leq \frac{n(n - 1)2^{\binom{n-1}{2}}}{2^{\binom{n}{2}}} = \frac{n(n - 1)}{2^{n-1}}.$$

Since clearly $\lim_{n \rightarrow \infty} \frac{n(n-1)}{2^{n-1}} = 0$, (31) is true. \square

Note that properties $\varphi_{\text{none}}(x)$ and $\varphi_{\text{all}}(x)$ can be shown to have typical degree 0 also by a direct computation, without much effort, without the help of Lemma 15. However such a direct proof for $\varphi_{\text{one}}(x)$ seems infeasible, because of the vast complexity of the required computations. Moreover the method of proof for $\varphi_{\text{one}}(x)$ used in the last proposition can be easily generalized in order to apply to the property “ x is adjacent to exactly k nodes”, for every $k \geq 1$.

Proposition 32. For each $k \geq 1$, let $\varphi_k(x)$ be the property “ x is adjacent to exactly k nodes”. Then $d(\varphi_k : \text{typ}) = 0$.

Proof. By Lemma 15(i), it suffices to show that $\mu(\varphi_k^{(1)}) = 0$. We argue as in the proof of 31 concerning $\varphi_{\text{one}}(x)$. Given a graph $G = \langle A, E \rangle$ with $|A| = n$ which satisfies $\varphi_k^{(1)}$, let $a \in A$ witness $\varphi_k^{(1)}$. Then a is adjacent exactly to k elements of $A \setminus \{a\}$. Since there are $\binom{n-1}{k}$ many k -element subsets of $A \setminus \{a\}$, there exist at most $\binom{n-1}{k} 2^{\binom{n-1}{2}}$ graphs on A in which a witnesses $\varphi_k^{(1)}$. Therefore, $|\text{Mod}_n(\varphi_k^{(1)})| \leq \binom{n-1}{k} 2^{\binom{n-1}{2}}$, and hence

$$\mu_n(\varphi_k^{(1)}) = \frac{|\text{Mod}_n(\varphi_k^{(1)})|}{|\mathbf{S}_n(L)|} \leq \frac{\binom{n-1}{k} 2^{\binom{n-1}{2}}}{2^{\binom{n}{2}}} = \frac{\binom{n-1}{k}}{2^{n-1}}.$$

In $\frac{\binom{n-1}{k}}{2^{n-1}}$ the nominator is a polynomial in n of degree $k + 1$, so by l' Hospital rule $\frac{\binom{n-1}{k}}{2^{n-1}} \xrightarrow{n} 0$. \square

Corollary 33. All properties $\varphi_{\text{none}}(x)$, $\varphi_{\text{all}}(x)$ and $\varphi_k(x)$, for $k \geq 1$, are regular.

Proof. We showed in the proofs of Propositions 31 and 32 that for each of the aforementioned properties $\varphi(x)$, $\mu(\varphi^{(1)}) = 0$. This condition implies, according to Lemma 15(ii), that $\varphi(x)$ is regular. \square

We close this section and the paper with some information offered by the reviewer about the previous results. Specifically, she or he kindly pointed out to me that there is an alternative, more “logical”, proof of the zero typicality degree of the properties $\varphi_{\text{none}}(x)$, $\varphi_{\text{all}}(x)$ and $\varphi_k(x)$, for $k \geq 1$, compared to the combinatorial proof we gave above. Namely, the negations of these properties follow from the random theory of undirected graphs which is axiomatized with so-called parametric sentences. One can consult [2, pp. 75–76] for details on this notion and the random theory of undirected graphs.

Acknowledgements. I would like to thank the anonymous reviewer for her or his thorough and very detailed checking which has led to some important improvements of the article.

References

- [1] T. M. Apostol. *Calculus*. Vol. 1. 2nd edition. New York: John Wiley & Sons, Inc., 1967.
- [2] H.-D. Ebbinghaus & J. Flum. *Finite Model Theory*. 1st edition. Perspectives in Mathematical Logic. Springer, 1995.
- [3] R. Fagin. Probabilities on finite models. *J. Symb. Log.* 41:1 (1976), pp. 50–58. DOI: 10.2307/2272945.
- [4] Y. V. Glebskii, D. I. Kogan, M. I. Liogon’kii & V. A. Talanov. Range and degree of realizability of formulas in the restricted predicate calculus. *Cybernetics* 5 (1969), pp. 142–154. DOI: 10.1007/BF01071084.
- [5] J. M. Harris, J. L. Hirst & M. J. Mossinghoff. *Combinatorics and Graph Theory*. 2nd edition. Undergraduate Texts in Mathematics. Springer, 2008.
- [6] B. C. Rennie & A. J. Dobson. On Stirling numbers of the second kind. *J. Comb. Theory* 7:2 (1969), pp. 116–121. DOI: 10.1016/S0021-9800(69)80045-1.
- [7] B. Russell. *My Philosophical Development*. Revised edition. Routledge, 1995.
- [8] R. Stanley. *Enumerative Combinatorics*. Vol. 1. 2nd edition. Cambridge University Press, 2012.
- [9] A. Tzouvaras. Russell’s typicality as another randomness notion. *Math. Log. Q.* 66:3 (2020), pp. 355–365. DOI: 10.1002/malq.202000038.
- [10] A. Tzouvaras. Typicality à la Russell in set theory. *Notre Dame J. Formal Logic* 63:2 (2022), pp. 185–196. DOI: 10.1215/00294527-2022-0011.